

© RUT Printer and Publisher

Print & Online, Open Access, Research Journal Available on <http://jbsd.in>

ISSN: 2229-3469 (Print); ISSN: 2231-024X (Online)

Research Article



Use of biopotentials of plant extracts of medicinal importance against pathogenic fungi *Fusarium oxysporum*.

Kirankumar R. Khandare and Rajendrabhai D. Vasait*

Mycology and Plant pathology Laboratory, Department of Botany, K.A.A.N.M.S. Arts, Comm. and Science College Satana, Dist. Nashik (M.S.) India. (drkhandarekr@gmail.com)

*Department of Microbiology, K.A.A.N.M.S. Arts, Comm. and Science College Satana, Dist. Nashik (M.S.) India. (rvasait@yahoo.com)

Article Info

Received: 02-10-2016,

Revised: 18-11-2016,

Accepted: 22-11-2016

Keywords:

Fusarium, Indian plants, PCE, Wilt.

Abstract

Stem, leaf, seed and root extract of ten Indian plants (*Accacia catechu* Willd, *Accacia arabicae* Willd, *Carissa congesta* Wight, *Cassia fistula* (L.), *Cuscuta reflexa* Roxb., *Jatropha curcas* L., *Moringa oleifera* Lam., *Syzygium cumini*, *Terminalia arjuna* Roxb., *Vitex negundo* L.) of medicinal importance from India significantly manage the *Fusarium* wilt of Tomato (*Lycopersicon esculantum*) caused by *Fusarium oxysporum* sp. *Lycopersicon*. Root extract of *Accacia catechu* control the growth of test fungi at 25% concentration (23.66), followed by leaf extract of *Jatropha curcas* at 25%, showed 33.33 cm growth of test fungi in treated plates, remaining extracts of plant parts inhibited the growth at 25% with less than 45cm growth. All extracts showed 100% PCE in 50% concentration against test fungi, while in 25% concentration it showed more than 99.8% PCE in several plant extracts.

INTRODUCTION

The use of medicinal plant materials for the inhibition of fungal diseases is an old practice in many countries and is still offers an enormous potential source of antifungal agent (Usharani and Chitra, 2014). Many phytochemical pesticides exhibiting broad spectrum of activity against pest and diseases have long been considered as attractive alternative to synthetic chemical pesticides as they are biodegradable, target specific and pose no or less hazard to the environment or to human health. (Walia *et al.*, 2014). Plants have been a rich source of medicines because the produce a bioactive molecules, most of which probably evolved as chemical defenses against predation or infection (Anitha *et al.*, 2016). Irrational use of chemicals pesticides pose dangerous to overall ecosystem these chemicals combat various phytopathogens which are inhibited various crop diseases, but on the other hand they have their adverse effect on human health and environment. Pesticides of chemical

origin pollutes water, air soil (Manuel *et al.*, 2008). Medicinal plants have played an essential role in the development of human culture. Medicinal plants are the resources of various medicinally important constituents, chemical ingredients (Saleh, 2015).

The testing of the efficacy of such potential plant based sources for antifungal activity could an important step towards the assessment of the degree of variability among the diverse natural flora (Manoorkar and Gachande, 2014). As plants contains various chemicals which showed antifungal, antibacterial activities, Therefore the present study was undertaken to manage fungal pathogen *Fusarium oxysporum* causing disease to a Tomato (*Lycopersicon esculantum*) through root, seed and leaf extracts of ten Indian plants of medicinal important (*Accacia catechu* Willd, *Accacia arabicae* Willd, *Carissa congesta* Wight, *Cassia fistula* (L.), *Cuscuta reflexa* Roxb., *Jatropha curcas* L., *Moringao leifera* Lam., *Syzygium cumini*, *Terminalia arjuna* Roxb., *Vitex negundo* L.).

MATERIALS AND METHODS

Fresh plant materials viz. leaves, stem, seeds and roots were collected from different parts of Maharashtra, India. These all ten plants (*Accacia catechu* Willd, *Accacia arabicae* Willd, *Carissa congesta* Wight, *Cassia fistula*(L.), *Cuscuta reflexa* Roxb., *Jatropha curcas* L., *Moringa oleifera* Lam., *Syzygium cumini*, *Terminalia arjuna* Roxb., *Vitex negundo* L.) have their own traditionally medicinal importance (Table No.1) collected plant materials were thoroughly washed with tap water and then followed by distilled water and kept in dark in filter papers at room temperature till completely dry. Each plant material was individually pulverized to obtained dry powder. For each plant 25 gm powder was taken. Extract of each plant material of every plant was prepared in ethanol and distilled water and serve as stock extract, the toxicity of extract was determined against pathogenic fungi, *Fusarium*

oxysporum following food poisoned technique (Mishra and Tiwari, 1992)

Different concentrations of extracts were prepared (10%, 25%, 50% and 100%) supplemented with Czepox Dox agar medium, four concentrations of each plant material of every plant were made each concentration with three replications and control were kept. These petriplates then inoculated with six mm disc of fungal mycelium, which is obtained from seven day old culture of fungal pathogen. These inoculated plates kept upside down and inoculated in BOD incubator at $28^{\circ}\text{C} \pm 2^{\circ}\text{C}$. Radial growths of fungal colonies were measured at different intervals. Antifungal activities of extracts of different plant parts of all ten plants was studied, sporulation of test fungi on different extracts was calculated by following equation.

$$\text{Spore per ml} = \frac{\text{No. of spores observed per microscopic field}}{12.5} \times \frac{40,000}{1}$$

The percentage control efficacy is also studied, which was calculated by following formula

$$\text{PCE} = 100 \left(1 - \frac{X}{Y} \right)$$

Where,

X = The diameter of the mycelia growth on extract treated host.

Y = The diameter of the mycelia growth on untreated host (Control)

RESULTS AND DISCUSSION

Leaves, stem, seed and root extracts of different ten plants tested against test fungi viz. *Fusarium oxysporum* significantly inhibited radial growth of fungal pathogen at various concentrations, as shown in Table 2. At 50% concentration extracts of different plant parts of ten plants significantly inhibited the test fungi *Fusarium oxysporum*, root extract of *Accacia catechu* control the growth of test fungi at 25% concentration (23.66), followed by leaf extract of *Jatropha curcas* at 25%, showed 33.33 cm growth of test fungi in treated plates, remaining extracts of plant parts inhibited the growth at 25% with less than 45cm growth (table 2).

In vivo percentage control efficacy (PCE) of all plant part extracts of ten plants was studied against test fungi. All extracts showed 100% PCE in 50% concentration against test fungi, while in 25% concentration it showed more than 99.8% PCE in several plant extracts (table 4) like *Accacia catechu*, *Crissa congesta*, *Jatropha curcas*, *Moringa oleifera* and *Vitex negundo*. Aqueous and ethanolic leaf extracts of *Calotropis procera* showed antifungal activities against dominant storage seed born fungi *Curularia lunata*, *Alternaria alternate*, *Rhizoctonia solani* *Fusarium solani*, *Penicillium chrysogenum*, *Aspergillus niger*, *A. flavus*, *A. terreus*, *A. fumigates*, and *Rhizopus sp.*(Manoorkar *et al*, 2015).

Table No. 1: Medicinally importance Indian plants.

Sr. No.	Botanical Name of plant	Family	Traditional use
1	<i>Accacia catechu</i> Willd	Mimosaceae	Used in the treatment of diarrhea and throat infections.
2	<i>Accacia arabicae</i> Willd	Mimosaceae	Skin wounds, such as burns, cuts or leprosy, digestive, gonorrhea, cough, dysentery, colds etc.
3	<i>Carissa congesta</i> Wight	Apocynaceae	Stomachic, antidiarrhoeal, vermifuge, antianthelmintic, astringent and antiascorbutic
4	<i>Cassia fistula</i> (L.)	Caesalpiniaceae	Anti-inflammatory, antioxidant, wound healing, laxative, purgative, useful on leucoderma, diabetes and antibacterial.
5	<i>Cuscuta reflexa</i> Roxb.	Convolvulaceae	Seeds are carminative and anthelmintic, plant used externally against itch, internally in protected fevers.
6	<i>Jatropha curcas</i> L.	Ephorbiaceae	Human and veterinary ailment, latex is disinfectant, anti cancerous, used in skin disease, piles, malaria, rheumatic and muscular pain.
7	<i>Moringa oleifera</i> Lam	Moringaceae	Antioxidant, antispasmodic, antiulcer anti-inflammatory, diuretic wound healing.
8	<i>Syzygium cumini</i>	Myrtaceae	Fruit used in diabetes, bark is anti-inflammatory, digestive ailments, leaves are antioxidants, used in blood pressure.
9	<i>Terminalia arjuna</i> Roxb.	Combretaceae	Antioxidant, anti mutagenic, cardiac tonic, anti cancerous.
10	<i>Vitex negundo</i> L.	Verbinaceae	Muscle relaxant, pain relieving, anti mosquito, anti anxiety, anti asthma

Table: 2 Antifungal activities plant extracts against *Fusarium oxysporum*

Sr. No.	Botanical extracts	Plant part used	Conc. of extract in%	Growth of <i>Fusarium oxysporum</i> in mm
1	<i>Accacia catechu</i> Willd	Root	10	55.33*
			25	23.66
			50	00.00
2	<i>Accacia arabicae</i> Willd	Seed	10	64.33
			25	43.66
			50	00.00
3	<i>Carissa congesta</i> Wight	Leaf	10	58.66
			25	39.33
			50	00.00
4	<i>Cassia fistula</i> (L.)	Seed	10	59.66
			25	41.66
			50	00.00
5	<i>Cuscuta reflexa</i> Roxb.	Stem	10	61.33
			25	45.33
			50	00.00
6	<i>Jatropha curcas</i> L.	Leaf	10	52.00
			25	33.33
			50	00.00
7	<i>Moringa oleifera</i> Lam	Leaf	10	59.00
			25	42.33
			50	00.00
8	<i>Syzygium cumini</i>	Seed	10	67.33
			25	45.66
			50	00.00
9	<i>Terminalia arjuna</i> Roxb.	Stem	10	62.33
			25	41.66
			50	00.00
10	<i>Vitex negundo</i> L.	Leaf	10	59.33
			25	36.33
			50	00.00
			Control	90.00

*Radial growth is the mean of triplicat

Table No 3: Sporulation of *Fusarium oxysporum* against different extracts.

Sr. No.	Botanical extracts	Plant part used	Conc. of extract in%	Sporulation of <i>Fusarium oxysporum</i> in mm	
				No. of spores per microscopic field*	No. of spores per ml suspension
1	<i>Accacia catechu</i> Willd	Root	10	4.0	12,800
			25	1.0	3,200
			50	0.0	00.00
2	<i>Accacia arabicae</i> Willd	Seed	10	5.0	16,00
			25	0.0	00.00
			50	0.0	00.00
3	<i>Carissa congesta</i> Wight	Leaf	10	7.0	22,400
			25	2.0	6,400
			50	0.0	00.00
4	<i>Cassia fistula</i> (L.)	Seed	10	5.0	16000
			25	1.0	3,200
			50	0.0	00.00
5	<i>Cuscuta reflexa</i> Roxb.	Stem	10	6.0	19,200
			25	3.0	9,600
			50	0.0	00.00
6	<i>Jatropha curcas</i> L.	Leaf	10	3.0	9,600
			25	1.0	3,200
			50	0.0	00.00
7	<i>Moringa oleifera</i> Lam	Leaf	10	7.0	22,400
			25	4.0	12,800
			50	0.0	00.00
8	<i>Syzygium cumini</i>	Seed	10	8.0	25,600
			25	4.0	12,800
			50	0.0	00.00
9	<i>Terminalia arjuna</i> Roxb.	Stem	10	6.0	19,200
			25	4.0	12,800
			50	0.0	00.00
10	<i>Vitex negundo</i> L.	Leaf	10	5.0	16,00
			25	3.0	9,600
			50	0.0	00.00
			Control	14.6	46,720

*No. of spores per microscopic field in 10X

According to Farzana *et al.*, (2014) The antifungal activity of ethanol and acetone extracts of leaves of nine medicinal plants i.e. *Piper betel*, *Lowsoniainermis*, *Psidiumuajava*, *Carissa papaya*, *Mimosa pudica*, *Catheranthusroseus*, *Moringa oleifera*, *Adhatoda vasica*, and *Andrographis paniculata* against *Fusarium oxysporum*, the causal organism of *Fusarium* wilt of Tomato was assessed 100%. Antifungal activity of different concentrations (2, 4, 6, and 8% w/v) of leaf extracts of wheat (*Triticum aestivum* L) Maize (*Zea mays* L) Sunflower (*Helianthus annus* L) Chilies (*Capsicum annum*), Onion (*Allium cepa*), and Marogold

(*Tagatus erectus* L) successfully inhibit corn rot disease of *Gladiolus* caused by *Fusarium oxysporum* (Tariq, 2008). Bhalodia *et al.*, 2011, tested *Cassia fistula* against two gram positive bacteria and three fungal pathogens i.e. *Aspergillus niger*, *Aspergillus clavatus* and *Candida albicans*, the result shows remarkable inhibition of the fungal growth. Methanol leaf extract of *Leucas zylanica* and *L. aspera*. Antifungal susceptibilities of certain clinically isolated dermatophytes to methanol extracts of *Leucas aspera* and *Leucas zeylanica* leaves were performed using agar well diffusion method.

Table No. 4: In vivo Percentage Control Efficacy (PCE) of the different extracts against test fungi

Sr. No.	Botanical extracts	Plant part used	Conc. of extract in%	<i>Fusarium oxysporum</i>	
				Growth in mm	PCE
1	<i>Accacia catechu</i> Willd	Root	10	17.0	99.6
			25	8.0	99.8
			50	0.0	100
2	<i>Accacia arabicae</i> Willd	Seed	10	19.0	99.5
			25	11.0	99.7
			50	0.0	100
3	<i>Carissa congesta</i> Wight	Leaf	10	18.0	99.6
			25	9.0	99.8
			50	0.0	100
4	<i>Cassia fistula</i> (L.)	Seed	10	20.0	99.5
			25	12.0	99.7
			50	0.0	100
5	<i>Cuscuta reflexa</i> Roxb.	Stem	10	21.0	99.5
			25	12.0	99.7
			50	0.0	100
6	<i>Jatropha curcas</i> L.	Leaf	10	15.0	99.6
			25	9.0	99.8
			50	0.0	100
7	<i>Moringa oleifera</i> Lam	Leaf	10	20.0	99.5
			25	9.0	99.8
			50	0.0	100
8	<i>Syzygium cumini</i>	Seed	10	22.0	99.5
			25	12.0	99.7
			50	0.0	100
9	<i>Terminalia arjuna</i> Roxb.	Stem	10	21.0	99.5
			25	12.0	99.7
			50	0.0	100
10	<i>Vitex negundo</i> L.	Leaf	10	20.0	99.5
			25	10.0	99.8
			50	0.0	100
			Control	42.0	00

The result obtained shows that all the extracts expressed remarkable antifungal activity with zone of inhibition ranging from 5 to 10 mm (Babu *et al.*, 2016). According to Gavande *et al.*, (2014) The antifungal potential seeds of *Aegle marmelos* tested against antraconose causing plant fungal pathogen *Colletotrichum acutatum*, fungal strain of *Metarhiziumani sopliae*, *Trichoderma aharzianum* and *Penicillium* sp., both ethanolic and petroleum ether extracts showed antifungal activities. Rathod *et al.*, (2015) used water and acetone extracts against fungus *Candida albicans* showed fungicidal properties, the presence of phytochemicals may be responsible for these activity, they found that acetone extracts are more efficient than that of aqueous extracts. Leaf extracts of ten plants (*Annona squamosa*, *Azadirachta*

indica, *Adhatoda vasica*, *Ocimum sanctum* , *Polyalthia longifolia*, *Tridax procumbens* , *Curcuma longa*, *Zingiber officinale*, *Allium cepa* and *Allium sativum*) found inhibitory against *Curvularia lunata*, *Phytophthora* sp., *Alternaria alternate*, *Fusarium oxysporum* *Asperillus niger* and *Rhizoctonia solani* (Swami and Alane, 2013).

According to Mogle and Maske, (2012) The effects of leaf extracts of *Argemone mexicana* L., *Semecarpus anacardium* L., *Cassia fistula* L., *Tephrosia purpurea* (L.) Pers., found useful for the control of *Collectotrichum destructivum* on seeds of cowpea. The seeds were soaked in sterile distilled water extract (10, 20 and 30%, w/v) of the leaves for 5, 10 and 15 h. All these plant extracts had significant inhibitory growth effect on the fungal pathogen.

Argemone mexicana extract was more effective followed by *Semecarpus anacardium*, *Cassia fistula* and *Tephrosia purpurea* plant extracts. Petroleum ether, hexane, chloroform, acetone and methanol extracts of leaf, stem and tuber of *Corallocarpus epigaeus* were investigated for antifungal activity against *Candida albicans*, *C. tropicalis*, *Aspergillus niger*, *A. flavus* and *A. versicolor* by disc diffusion method. Methanol extract of *C. epigaeus* tuber exhibited maximum activity against most of the tested fungi. The petroleum ether and hexane extracts obtained from *Corallocarpus epigaeus* stem was found to be active only against *A. niger*, *A. flavus* and *A. versicolor*. All the crude extracts exhibited activity against *A. niger* and *A. flavus* (Priyavardhini et al., 2012).

REFERENCES

- Anitha C, Soumaya JD, Ajitha RK., Remaya MP, Sreja JR and Suvachan A, 2016.** Augmentative biopotential efficiency of leaf extract on the tropical medicinal plant *Cardiospermum helicacabum*. *World Journal of Parasitology and Pharmaceutical Science*. **5**(9):1557-1568.
- Babu M, Sheik N, Mohamed K, Anand JD and P Saravanan, 2016.** In-vitro antifungal activity of leaf extracts of *Leucas aspera* and *Leucas zylanica*. *International Journal of Pharmaceutical Sciences and Research*, **7**(2):752-756.
- Farzana AN, Ismat A and Shanim S, 2014.** Antifungal activity of selected medicinal plant extracts on *Fusarium oxysporum schlechtthe*, causal organism of *Fusarium* wilt diseases in Tomato. *Biomedical and Life Sciences*. **5** (18):2665-2671.
- Gavanda AE, Gaikwad NS and Bale SR, 2014.** Selective inhibitory in vitro activity of *Aegle marmelos* (L.) extracts on plant fungal pathogen *Colletotrichum acutatum*. *Bioscience Discovery*. **5**(1):55- 59.
- Manoorkar VB, Mandge SV and Gachande BD, 2015.** Antifungal Activity of Leaf and Latex Extracts of *Calotropis procera* (Ait.) against Dominant Seed-Borne Storage Fungi of Some Oil Seeds. *Bioscience Discovery*. **6**(1):22-26.
- Manoorkar VB and Gachande BD, 2014.** Evaluation of antifungal activity of some medicinal plant extracts against some storage seed -borne fungi of Groundnut. *Science Research Reporter*, **4**(1):67-70.
- Manuel AE, Eugenio LP, Elena MC, Jesus SG, and Jaun CM, 2008.** The mobility and degradation of pesticides in soils and the pollution of groundwater resources, *Agriculture, Ecosystem and Environment*. **123**(4):247-260.
- Mishra M and Tiwari SN, 1992.** Toxicity of *Polyalthia longifolia* against fungal pathogens of rice. *Indian Phytopathology*. **45** (1): 192-198.
- Mogle UP and Maske SR, 2012.** Effect of some plant extract against seed borne infection of *Collectotrichum destructivum* on *Vigna unguiculata*. *Bioscience Discovery* **3**(2): 266 -269.
- Nayan RB, and VJ Shukla, 2011.** Antibacterial and antifungal activities from leaf extracts of *Cassia fistula* L.: An Ethnomedicinal plant. *J. of Advanced Pharm. Technology and Research*. **2**(2):104-109.
- Rathod M, Das N and Dhate DA, 2015.** Antifungal activity of two medicinal plants against fungus *Candida albicans*. *International Journal of Pharma and Bioscience*. **6**(4):701-706.
- Saleh H, Azizollah J, Ahmadreza H Raham A, 2015.** The applications of medicinal plants in traditional and modern medicine. *International Journal of Clinical Medicine*. **6**:635-542.
- Swami CS and Alane SK, 2013.** Efficacy of some botanicals against seed - borne fungi of green gram (*Phaseolus aureus roxb.*). *Bioscience Discovery*. **4**(1):107-110.
- Tariq R, Khan SN and Arshad J, 2008.** *Mycopath*. **6** (1 and 2):13-15.
- Usharani S and Chitra M, 2014.** Biopotential activity of a rare medicinal plant – *Wattaka kavolubilis* leaf. *International Journal of Pharm.Tech Research*. **6**(5):1584.
- Vasantha P, Tresina S and Veerabhau RM, 2012.** Antifungal activity of *Kedrostis foetidissima* (Jacq.) Cogn. *International Journal of PharmTech Research*. **4**(1):44-48.
- Walia S, Saha S and Virendra SR, 2014.** Phytochemical pesticides, Advances in plant Biopesticides. *Springer India*. 295-322.

How to Cite this Article:

Kirankumar R. Khandare and Rajendrabhai D. Vasait, 2017. Use of biopotentials of plant extracts of medicinal importance against pathogenic fungi *Fusarium oxysporum*. *Bioscience Discovery*, **8**(1):87-92.